arXiv:1607.01343v1 [math.CA] 5 Jul 2016

In memory of my Grandmother

ON THE SHARPNESS OF ONE INEQUALITY OF DIFFERENT METRICS FOR ALGEBRAIC POLYNOMIALS

Roman A. Veprintsev

Abstract. We prove that the previously established inequality of different metrics for algebraic polynomials is sharp in the sense of order.

Key words and phrases: inequality of different metrics, algebraic polynomials, generalized Jacobi weight, orthogonal polynomials, generalized Gegenbauer polynomials

MSC 2010: 33C45, 33C47, 41A17

1. Introduction and preliminaries

In this section, we give some notation used in the article. Here, we also introduce some classes of orthogonal polynomials on $[-1,1]$, including the so-called generalized Gegenbauer polynomials. For a background and more details on the orthogonal polynomials, the reader is referred to [1–3, 5].

Consider the generalized Jacobi weight

$$\omega_{\alpha,\beta,\gamma}(x) = (1-x)^\alpha (1+x)^\beta |x|^\gamma, \qquad x \in [-1,1],$$

with α, β, $\gamma > -1$. Let $\{p_n^{(\alpha,\beta,\gamma)}\}_{n=0}^\infty$ denote the sequence of orthonormal polynomials with respect to the weight function $\omega_{\alpha,\beta,\gamma}$.

Given $1 \le p \le \infty$, we denote by $L_p(\omega_{\alpha,\beta,\gamma})$ the space of complex-valued Lebesgue measurable functions f on $[-1,1]$ with finite norm

$$\|f\|_{L_p(\omega_{\alpha,\beta,\gamma})} = \Big(\int_{-1}^1 |f(x)|^p\, \omega_{\alpha,\beta,\gamma}(x)\, dx\Big)^{1/p}, \qquad 1 \le p < \infty,$$

$$\|f\|_{L_\infty(\omega_{\alpha,\beta,\gamma})} = \operatorname*{ess\,sup}_{x\in[-1,1]} |f(x)|.$$

The Jacobi polynomials, denoted by $P_n^{(\alpha,\beta)}(\cdot)$, where $n = 0, 1, \ldots$, are orthogonal with respect to the Jacobi weight function $w_{\alpha,\beta}(x) = (1-x)^\alpha (1+x)^\beta$ on $[-1,1]$, namely,

$$\int_{-1}^1 P_n^{(\alpha,\beta)}(x)\, P_m^{(\alpha,\beta)}(x)\, w_{\alpha,\beta}(x)\, dt = \begin{cases} \dfrac{2^{\alpha+\beta+1}\Gamma(n+\alpha+1)\Gamma(n+\beta+1)}{(2n+\alpha+\beta+1)\Gamma(n+1)\Gamma(n+\alpha+\beta+1)}, & n = m, \\ 0, & n \ne m. \end{cases}$$

Here, as usual, Γ is the gamma function. Note that $w_{\alpha,\beta} = \omega_{\alpha,\beta,0}$.

For $\lambda > -\frac{1}{2}$, $\mu \geq 0$, and $n = 0, 1, \ldots$, the orthonormal generalized Gegenbauer polynomials $\widetilde{C}_n^{(\lambda,\mu)}(\cdot)$ are defined by

$$\begin{aligned}
&\widetilde{C}_{2n}^{(\lambda,\mu)}(x) = \widetilde{a}_{2n}^{(\lambda,\mu)} P_n^{(\lambda-1/2,\mu-1/2)}(2x^2-1),\\
&\widetilde{a}_{2n}^{(\lambda,\mu)} = \left(\frac{(2n+\lambda+\mu)\Gamma(n+1)\Gamma(n+\lambda+\mu)}{\Gamma(n+\lambda+\frac{1}{2})\Gamma(n+\mu+\frac{1}{2})}\right)^{1/2},\\
&\widetilde{C}_{2n+1}^{(\lambda,\mu)}(x) = \widetilde{a}_{2n+1}^{(\lambda,\mu)}\, x P_n^{(\lambda-1/2,\mu+1/2)}(2x^2-1),\\
&\widetilde{a}_{2n+1}^{(\lambda,\mu)} = \left(\frac{(2n+\lambda+\mu+1)\Gamma(n+1)\Gamma(n+\lambda+\mu+1)}{\Gamma(n+\lambda+\frac{1}{2})\Gamma(n+\mu+\frac{3}{2})}\right)^{1/2}.
\end{aligned} \tag{1}$$

They are orthonormal with respect to the weight function

$$v_{\lambda,\mu}(x) = (1-x^2)^{\lambda-1/2}|x|^{2\mu}, \quad x \in [-1,1].$$

Note that $v_{\lambda,\mu} = \omega_{\lambda-1/2,\lambda-1/2,2\mu}$ and

$$\widetilde{C}_n^{(\lambda,\mu)} = p_n^{(\lambda-1/2,\lambda-1/2,2\mu)}. \tag{2}$$

The generalized Gegenbauer polynomials play an important role in Dunkl harmonic analysis (see, for example, [2, 3]). So, the study of these polynomials and their applications is very natural.

Define the uniform norm of a continuous function f on $[-1,1]$ by $\|f\|_\infty = \max\limits_{-1\leq x\leq 1} |f(x)|$. The maximum of two real numbers x and y is denoted by $\max(x,y)$.

Throughout the paper we use the following asymptotic notation: $f(n) \lesssim g(n)$, $n \to \infty$, or equivalently $g(n) \gtrsim f(n)$, $n \to \infty$, means that there exist a positive constant C and a positive integer n_0 such that $0 \leq f(n) \leq Cg(n)$ for all $n \geq n_0$ (asymptotic upper bound); if there exist positive constants C_1, C_2, and a positive integer n_0 such that $0 \leq C_1 g(n) \leq f(n) \leq C_2 g(n)$ for all $n \geq n_0$, then we write $f(n) \asymp g(n)$, $n \to \infty$ (asymptotic tight bound).

To simplify the writing, we will omit "$n \to \infty$" in the asymptotic notation.

It follows directly from Stirling's asymptotic formula that

$$\frac{\Gamma(n+\alpha)}{\Gamma(n+\beta)} \asymp n^{\alpha-\beta} \tag{3}$$

for arbitrary real numbers α and β.

It is known [1, p. 350] that

$$\|P_n^{(\alpha,\beta)}\|_\infty \asymp \begin{cases} n^{\max(\alpha,\beta)} & \text{if } \max(\alpha,\beta) \geq -1/2,\\ n^{-1/2} & \text{if } \max(\alpha,\beta) < -1/2. \end{cases} \tag{4}$$

The aim of the paper is to prove that the established in [7] inequality of different metrics for algebraic polynomials is sharp in the sense of order.

2. Main result

In [7], we prove the following result.

Theorem 1. *Let $\alpha \geq \beta \geq -\frac{1}{2}$, $\mu \geq 0$, $1 \leq p < q \leq \infty$. If P_n is an algebraic polynomial of degree n, then*

$$\|P_n\|_{L_q(\omega_{\alpha,\beta,\mu})} \lesssim n^{\max(2(\alpha+1),\mu+1)\left(\frac{1}{p}-\frac{1}{q}\right)} \|P_n\|_{L_p(\omega_{\alpha,\beta,\mu})}, \tag{5}$$

i.e.,

$$\sup_{P_n} \frac{\|P_n\|_{L_q(\omega_{\alpha,\beta,\mu})}}{\|P_n\|_{L_p(\omega_{\alpha,\beta,\mu})}} \lesssim n^{\max(2(\alpha+1),\mu+1)\left(\frac{1}{p}-\frac{1}{q}\right)},$$

where the supremum is taken over all polynomials P_n of degree n.

Now we can formulate the main result.

Theorem 2. *Let $\alpha \geq \beta \geq -\frac{1}{2}$, $\mu \geq 0$, $1 \leq p < q \leq \infty$, and $\nu \in \left(0, 1-\frac{1}{q}\right)$. Then*

$$\sup_{P_n} \frac{\|P_n\|_{L_q(\omega_{\alpha,\beta,\mu})}}{\|P_n\|_{L_p(\omega_{\alpha,\beta,\mu})}} \gtrsim \begin{cases} n^{\max(2(\alpha+1),\mu+1)\left(\frac{1}{p}-\frac{1}{q}\right)} & \text{if } 1 < p < q \leq \infty, \\ n^{\max(2(\alpha+1),\mu+1)\left(1-\frac{1}{q}\right)-\nu} & \text{if } p = 1,\ 1 < q \leq \infty, \end{cases}$$

where the supremum is taken over all polynomials P_n of degree n.

Thus, the inequality (5) of different metrics for algebraic polynomials is precise in order.

Corollary 1. *Let $\alpha \geq \beta \geq -\frac{1}{2}$, $\mu \geq 0$, $1 < p < q \leq \infty$. Then*

$$\sup_{P_n} \frac{\|P_n\|_{L_q(\omega_{\alpha,\beta,\mu})}}{\|P_n\|_{L_p(\omega_{\alpha,\beta,\mu})}} \asymp n^{\max(2(\alpha+1),\mu+1)\left(\frac{1}{p}-\frac{1}{q}\right)},$$

where the supremum is taken over all polynomials P_n of degree n.

This corollary immediately follows from Theorems 1 and 2.

3. Some auxiliary results

In this section, we present some results used in the proof of Theorem 2.
In [4] it was proved the following result.

Lemma 1. *Let α, $\beta \geq -1/2$, $\gamma \geq 0$, and $\widetilde{\alpha}$, $\widetilde{\beta}$, $\widetilde{\gamma} > -1$. Let $-1 < y_1 < 0 < y_2 < 1$. Then, for $1 \leq p < \infty$,*

$$\int_{y_2}^{1} (1-x)^{\widetilde{\alpha}} |p_n^{(\alpha,\beta,\gamma)}(x)|^p \, dx \asymp \begin{cases} 1 & \text{if } 2\widetilde{\alpha} > p\alpha - 2 + p/2, \\ \ln n & \text{if } 2\widetilde{\alpha} = p\alpha - 2 + p/2, \\ n^{p\alpha+p/2-2\widetilde{\alpha}-2} & \text{if } 2\widetilde{\alpha} < p\alpha - 2 + p/2, \end{cases}$$

$$\int_{y_1}^{y_2} |x|^{\widetilde{\gamma}} |p_n^{(\alpha,\beta,\gamma)}(x)|^p \, dx \asymp \begin{cases} 1 & \text{if } 2\widetilde{\gamma} > p\gamma - 2, \\ \ln n & \text{if } 2\widetilde{\gamma} = p\gamma - 2, \\ n^{p\gamma/2 - \widetilde{\gamma} - 1} & \text{if } 2\widetilde{\gamma} < p\gamma - 2, \end{cases}$$

$$\int_{-1}^{y_1} (1+x)^{\widetilde{\beta}} |p_n^{(\alpha,\beta,\gamma)}(x)|^p \, dx \asymp \begin{cases} 1 & \text{if } 2\widetilde{\beta} > p\beta - 2 + p/2, \\ \ln n & \text{if } 2\widetilde{\beta} = p\beta - 2 + p/2, \\ n^{p\beta + p/2 - 2\widetilde{\beta} - 2} & \text{if } 2\widetilde{\beta} < p\beta - 2 + p/2. \end{cases}$$

Using this lemma and (2), one can deduce the following two corollaries.

Corollary 2. *Let $\alpha \geq \beta \geq -\frac{1}{2}$, $\mu \geq 0$, and $1 < p < \infty$. Then*

$$\|p_n^{(2\alpha+\frac{3}{2}, 2\beta+\frac{3}{2}, 2\mu+2)}\|_{L_p(\omega_{\alpha,\beta,\mu})} \asymp n^{\max(2(\alpha+1),\mu+1)\left(1-\frac{1}{p}\right)}.$$

In particular,

$$\|\widetilde{C}_n^{(2(\alpha+1),\mu+1)}\|_{L_p(\omega_{\alpha,\beta,\mu})} \asymp n^{\max(2(\alpha+1),\mu+1)\left(1-\frac{1}{p}\right)}.$$

Corollary 3. *Let $\alpha \geq \beta \geq -\frac{1}{2}$, $\mu \geq 0$, $\nu \in \left(0, 1-\frac{1}{q}\right)$, and $1 < q < \infty$. Then*

$$\|p_n^{(2\alpha+\frac{3}{2}-\nu, 2\beta+\frac{3}{2}-\nu, 2\mu+2-2\nu)}\|_{L_1(\omega_{\alpha,\beta,\mu})} \asymp 1,$$
$$\|p_n^{(2\alpha+\frac{3}{2}-\nu, 2\beta+\frac{3}{2}-\nu, 2\mu+2-2\nu)}\|_{L_q(\omega_{\alpha,\beta,\mu})} \asymp n^{\max(2(\alpha+1),\mu+1)\left(1-\frac{1}{q}\right)-\nu}.$$

In particular,

$$\|\widetilde{C}_n^{(2(\alpha+1)-\nu,\mu+1-\nu)}\|_{L_1(\omega_{\alpha,\beta,\mu})} \asymp 1.$$

For $\lambda > -\frac{1}{2}$, $\mu > 0$, we have [6, Theorem 1]

$$\|\widetilde{C}_n^{(\lambda,\mu)}\|_\infty \asymp n^{\max(\lambda,\mu)}. \tag{6}$$

The above formula is sufficient for our purpose, but let us show that this is valid for $\mu = 0$.

Proposition 1. *Let $\lambda > -\frac{1}{2}$. Then*

$$\|\widetilde{C}_n^{(\lambda,0)}\|_\infty \asymp n^{\max(\lambda,0)}.$$

P r o o f. From (1) and [5, Theorem 4.1], we have

$$\begin{aligned} \widetilde{C}_{2n}^{(\lambda,0)}(x) &= \widetilde{a}_{2n}^{(\lambda,0)} \, P_n^{(\lambda-1/2,-1/2)}(2x^2-1) = \\ &= \widetilde{a}_{2n}^{(\lambda,0)} \, \frac{\Gamma(n+\lambda+\frac{1}{2})\,\Gamma(2n+1)}{\Gamma(2n+\lambda+\frac{1}{2})\,\Gamma(n+1)} \, P_{2n}^{(\lambda-1/2,\lambda-1/2)}(x), \\ \widetilde{C}_{2n+1}^{(\lambda,0)}(x) &= \widetilde{a}_{2n+1}^{(\lambda,0)} \, x P_n^{(\lambda-1/2,1/2)}(2x^2-1) = \\ &= \widetilde{a}_{2n+1}^{(\lambda,0)} \, \frac{\Gamma(n+\lambda+\frac{1}{2})\,\Gamma(2n+2)}{\Gamma(2n+\lambda+\frac{3}{2})\,\Gamma(n+1)} \, P_{2n+1}^{(\lambda-1/2,\lambda-1/2)}(x). \end{aligned}$$

By (1) and (3),

$$\widetilde{a}_{2n}^{(\lambda,0)} \asymp n^{1/2}, \qquad \widetilde{a}_{2n+1}^{(\lambda,0)} \asymp n^{1/2},$$
$$\frac{\Gamma(n+\lambda+\frac{1}{2})\,\Gamma(2n+1)}{\Gamma(2n+\lambda+\frac{1}{2})\,\Gamma(n+1)} \asymp 1, \qquad \frac{\Gamma(n+\lambda+\frac{1}{2})\,\Gamma(2n+2)}{\Gamma(2n+\lambda+\frac{3}{2})\,\Gamma(n+1)} \asymp 1.$$

Hence, considering (4), we get the desired asymptotic behavior

$$\|\widetilde{C}_n^{(\lambda,0)}\|_\infty \asymp n^{1/2}\|P_n^{(\lambda-1/2,\lambda-1/2)}\|_\infty \asymp$$
$$\asymp n^{1/2}\cdot\begin{cases} n^{\lambda-1/2} & \text{if } \lambda \geq 0, \\ n^{-1/2} & \text{if } -1/2 < \lambda < 0. \end{cases}$$

□

4. Proof of Theorem 2

Let us consider the following cases:

I) $1 < p < q < \infty$ and $1 < p < \infty$, $q = \infty$;

II) $p = 1$, $1 < q < \infty$ and $p = 1$, $q = \infty$.

Case I). By Corollary 2 and (6), we get the desired behavior

$$\sup_{P_n}\frac{\|P_n\|_{L_q(\omega_{\alpha,\beta,\mu})}}{\|P_n\|_{L_p(\omega_{\alpha,\beta,\mu})}} \geq \frac{\|p_n^{(2\alpha+\frac{3}{2},2\beta+\frac{3}{2},2\mu+2)}\|_{L_q(\omega_{\alpha,\beta,\mu})}}{\|p_n^{(2\alpha+\frac{3}{2},2\beta+\frac{3}{2},2\mu+2)}\|_{L_p(\omega_{\alpha,\beta,\mu})}} \gtrsim$$
$$\gtrsim \frac{n^{\max(2(\alpha+1),\mu+1)\left(1-\frac{1}{q}\right)}}{n^{\max(2(\alpha+1),\mu+1)\left(1-\frac{1}{p}\right)}} = n^{\max(2(\alpha+1),\mu+1)\left(\frac{1}{p}-\frac{1}{q}\right)}$$

and

$$\sup_{P_n}\frac{\|P_n\|_\infty}{\|P_n\|_{L_p(\omega_{\alpha,\beta,\mu})}} \geq \frac{\|\widetilde{C}_n^{(2(\alpha+1),\mu+1)}\|_\infty}{\|\widetilde{C}_n^{(2(\alpha+1),\mu+1)}\|_{L_p(\omega_{\alpha,\beta,\mu})}} \gtrsim$$
$$\gtrsim \frac{n^{\max(2(\alpha+1),\mu+1)}}{n^{\max(2(\alpha+1),\mu+1)\left(1-\frac{1}{p}\right)}} = n^{\max(2(\alpha+1),\mu+1)\frac{1}{p}}.$$

Case II). Using Corollary 3 and (6), we obtain the following asymptotic estimates

$$\sup_{P_n}\frac{\|P_n\|_{L_q(\omega_{\alpha,\beta,\mu})}}{\|P_n\|_{L_1(\omega_{\alpha,\beta,\mu})}} \geq \frac{\|p_n^{(2\alpha+\frac{3}{2}-\nu,2\beta+\frac{3}{2}-\nu,2\mu+2-2\nu)}\|_{L_q(\omega_{\alpha,\beta,\mu})}}{\|p_n^{(2\alpha+\frac{3}{2}-\nu,2\beta+\frac{3}{2}-\nu,2\mu+2-2\nu)}\|_{L_1(\omega_{\alpha,\beta,\mu})}} \gtrsim$$
$$\gtrsim n^{\max(2(\alpha+1),\mu+1)\left(1-\frac{1}{q}\right)-\nu}$$

and

$$\sup_{P_n}\frac{\|P_n\|_\infty}{\|P_n\|_{L_1(\omega_{\alpha,\beta,\mu})}} \geq \frac{\|\widetilde{C}_n^{(2(\alpha+1)-\nu,\mu+1-\nu)}\|_\infty}{\|\widetilde{C}_n^{(2(\alpha+1)-\nu,\mu+1-\nu)}\|_{L_1(\omega_{\alpha,\beta,\mu})}} \gtrsim$$
$$\gtrsim n^{\max(2(\alpha+1),\mu+1)-\nu}.$$

REFERENCES

1. G. E. Andrews, R. Askey, and R. Roy, *Special Functions*, Encyclopedia of Mathematics and its Applications **71**, Cambridge University Press, Cambridge, 1999.
2. F. Dai and Y. Xu, *Approximation theory and harmonic analysis on spheres and balls*, Springer Monographs in Mathematics, Springer, 2013.
3. C. F. Dunkl and Y. Xu, *Orthogonal polynomials of several variables*, 2nd ed., Encyclopedia of Mathematics and its Applications **155**, Cambridge University Press, Cambridge, 2014.
4. B. Xh. Fejzullahu, On orthogonal expansions with respect to the generalized Jacobi weight, *Results. Math.* **63** (2013), 1177–1193.
5. G. Szegö, *Orthogonal polynomials*, 4th ed., American Mathematical Society Colloquium Publications **23**, American Mathematical Society, Providence, Rhode Island, 1975.
6. R. A. Veprintsev, On the asymptotic behavior of the maximum absolute value of generalized Gegenbauer polynomials, arXiv preprint 1602.01023 (2016).
7. R. A. Veprintsev, On an inequality of different metrics for algebraic polynomials, arXiv preprint 1606.06149 (2016).

Independent researcher, Uzlovaya, Russia
E-mail address: **veprintsevroma@gmail.com**